




Contents lists available at ScienceDirect

Ain Shams Engineering Journal

journal homepage: [www.sciencedirect.com](http://www.sciencedirect.com)

# A direct method for solving fractional order variational problems by hat basis functions

Osama H. Mohammed

Department of Mathematics and Computer Applications, College of Science, Al-Nahrain University, Baghdad, Iraq

## ARTICLE INFO

### Article history:

Received 27 April 2016

Revised 2 November 2016

Accepted 9 November 2016

Available online xxxx

### Keywords:

Fractional calculus

Fractional variational problems

Hat basis functions

Fractional Euler-Lagrange equations

## ABSTRACT

This paper presents a numerical technique for solving a class of fractional variational problems using a direct method based on operational matrix of generalized hat basis function. The fractional derivative is defined in the Caputo sense.

Minimization of such functional leads to a set of algebraic equations which are solved using an appropriate numerical technique. For the class of problems considered, the numerical solution can be obtained directly from the functional and there is no need to solve the fractional Euler-Lagrange equations.

Numerical examples are introduced combined with their approximate solutions and comparison with other numerical approach for conforming the accuracy and applicability of the proposed method.

© 2016 Ain Shams University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Fractional calculus is a field of applied mathematics that deals with derivative and integrals of arbitrary orders (including complex orders), and their applications in science, engineering, mathematics, economics, and other fields [1].

Fractional derivatives provide an excellent instrument for the description of memory and hereditary properties of various materials and processes. This is the main advantage of fractional derivatives in comparison with classical integer-order models, in which such effects are in fact neglected. The advantages of fractional derivatives become apparent in modeling mechanical and electrical properties of real materials, as well as in the description of rheological properties of rocks, and in many other fields [2].

Most fractional order differential equations do not have exact solutions, so approximate and numerical techniques must be used. Several numerical and approximate methods to solve fractional order differential equations have been given such as Adomian decomposition method [3], variational iteration method [4], homotopy analysis method [5], homotopy perturbation method [6], collocation method [7,8], wavelet method [9], finite element method [10] and spectral tau method [11].

Recently, the operational matrices of fractional derivatives and integrals have been derived for some types of orthogonal polynomials such as, the Legendre polynomials [12–16], Chebyshev

polynomials [17,18], Jacobi polynomials [19–23] and Laguerre polynomials [24] and used to solve several types of fractional order differential equations see [25,26].

A fractional calculus of variations problem is a subtopic of fractional calculus and it is a problem in which either the objective functional or the constraint equation or both contain at least one fractional derivative term [27].

This occurs naturally in many problems of physics, mechanics and engineering in order to provide more accurate models of physics phenomena (see [28,29]).

It is remarkable to refer that the starting point of fractional calculus of variation appear to be the Refs. [30,31], where Riewe developed the non concentrative Lagrangian, Hamiltonian, and other concepts of classical mechanics using fractional derivative.

After that numerous works have been dedicated to the fractional calculus of variation see for example [32–37].

In recent years, many analytical and numerical techniques have been used for obtaining the solution of fractional variational problems among them for instance [27,38–44], (see also [45–48]).

Babolian and Mordad [49] have used hat basis function for solving system of linear and nonlinear integral equations of the second kind by the hat functions. In [50], Tripathi et al. have used generalized hat basis functions to obtain approximate solutions of linear and nonlinear fractional differential equations. Heydari et al. [51] have used generalized hat basis functions for solving stochastic integro-differential equations. In [52] Heydari et al. used generalized hat basis functions for solving nonlinear stochastic integro-differential equations. In [53] Heydari et al. have used hat basis functions for solving a class of fractional optimal control problems.

Peer review under responsibility of Ain Shams University.

E-mail addresses: [uham@sci.nahrainuniv.edu.iq](mailto:uham@sci.nahrainuniv.edu.iq), [dr\\_usama79@yahoo.com](mailto:dr_usama79@yahoo.com)

<http://dx.doi.org/10.1016/j.asej.2016.11.006>

2090-4479/© 2016 Ain Shams University. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Mohammed OH. A direct method for solving fractional order variational problems by hat basis functions. Ain Shams Eng J (2016), <http://dx.doi.org/10.1016/j.asej.2016.11.006>

In this paper a generalized hat basis function will be used for solving a class of fractional variational problems.

This paper is organized as follows: In Section 2 some definitions of fractional order derivatives and integration are investigated, in Section 3 we describe the generalized hat basis functions and their properties. In Section 4 operational matrices of fractional and integer order integration for the generalized hat functions is given. In Section 5 the proposed method is described for solving fractional order variational problems by generalized hat basis functions. In Section 6 we introduce some illustrative examples to show the accuracy and efficiency of the proposed method. Finally a conclusion is drawn in Section 7.

## 2. Fractional order integration and derivative

There are various definitions of fractional integration and derivative such as Riemann-Liouville and Caputo definitions. Comparatively, the Caputo derivative has certain advantages when trying to model real-world phenomena with fractional differential equations.

**Definition 1.** The Riemann-Liouville fractional integral of order  $\alpha > 0$  is defined as follows:

$$I_x^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (t - \tau)^{\alpha-1} f(\tau) d\tau, \quad x > 0, \quad \alpha \in \mathbb{R}^+$$

where  $\Gamma(\alpha)$  is the Gamma function.

**Definition 2.** The Caputo fractional derivative of order  $\alpha > 0$  is defined as follows:

$${}_0^C D_x^\alpha f(x) = \begin{cases} \frac{1}{\Gamma(m-\alpha)} \int_0^x \frac{f^{(m)}(\tau)}{(x-\tau)^{\alpha+1-m}} d\tau & m-1 < \alpha < m \\ \frac{d^m}{dx^m} f(x) & \alpha = m \end{cases}$$

for  $\alpha > 0$ , we have the following properties of the Caputo fractional derivative:

- (i)  ${}_0^C D_x^\alpha (I_x^\alpha f(x)) = f(x)$ .
- (ii)  $I_x^\alpha ({}_0^C D_x^\alpha f(x)) = f(x) - \sum_{k=0}^{n-1} f^{(k)}(0^+) \frac{x^k}{k!}$ .
- (iii)  ${}_0^C D_x^\alpha (C) = 0, C \in \mathbb{R}$ .
- (iv)  ${}_0^C D_x^\alpha (x^\gamma) = \begin{cases} \frac{\Gamma(\gamma+1)}{\Gamma(\gamma-\alpha+1)} x^{\gamma-\alpha} & \gamma \in \{0, 1, 2, 3, \dots\}, \quad \gamma \geq [\alpha] \\ 0 & \gamma \in \{0, 1, 2, 3, \dots\}, \quad \gamma < [\alpha] \end{cases}$

where  $[\alpha]$  is the floor function of  $\alpha$ .

## 3. Generalized hat functions and their properties

Usually hat functions are defined on the interval  $[0, 1]$ . These are continuous functions with shape of hats, when plotted on two dimensional planes. In this section the generalized hat functions [50] are considered on the interval  $[0, T]$ . The interval  $[0, T]$  is divided into  $n$  subintervals  $[ih, (i+1)h]$ ,  $i = 0, 1, 2, \dots, n-1$ ; of equal lengths  $h$  where  $h = \frac{T}{n}$ . The generalized hat functions family of first  $(n+1)$  hat functions is defined as follows:

$$\psi_0(x) = \begin{cases} \frac{h-x}{h}, & 0 \leq x < h \\ 0, & \text{otherwise} \end{cases} \quad (1)$$

$$\psi_i(x) = \begin{cases} \frac{x-(i-1)h}{h}, & (i-1)h \leq x \leq ih \\ \frac{(i+1)h-x}{h}, & ih \leq x \leq (i+1)h, \quad i = 1, 2, \dots, n-1 \\ 0, & \text{otherwise} \end{cases} \quad (2)$$

$$\psi_n(x) = \begin{cases} \frac{x-(T-h)}{h}, & T-h \leq x \leq T \\ 0, & \text{otherwise} \end{cases} \quad (3)$$

These  $\psi_i(x)^s \in L^2[0, T]$  and are linearly independent. Using the definition of generalized hat functions, we have the following observations.

$$\psi_i(kh) = \begin{cases} 1, & i = k \\ 0, & i \neq k \end{cases} \quad (4)$$

and

$$\psi_i(x)\psi_j(x) = 0, \quad |i-j| \geq 2 \quad (5)$$

Any arbitrary function  $f \in L^2[0, T]$  is approximated in vector form as:

$$f(x) \cong \sum_{i=0}^{n-1} f_i \psi_i(x) = F_{n+1}^T \psi_{n+1}(x) = \psi_{n+1}^T F_{n+1} \quad (6)$$

where

$$F_{n+1} = [f_0, f_1, f_2, \dots, f_n]^T \quad (7)$$

and

$$\Psi_{n+1}(x) = [\psi_0(x), \psi_1(x), \psi_2(x), \dots, \psi_n(x)]^T \quad (8)$$

The importance aspect of using the generalized hat function in the approximation of function  $f(x)$ , lies in the fact that the coefficient  $f_i$  in Eq. (6), are given by:

$$f_i = f(ih), i = 0, 1, 2, \dots, n \quad (9)$$

Moreover if  $T = 1$  then from Eq.(5), we have

$$\int_0^1 \Psi_{n+1}(x) \Psi_{n+1}^T(x) dx = K_{n+1 \times n+1}$$

$$\text{where } K_{n+1 \times n+1} = \frac{h}{6} \begin{bmatrix} 2 & 1 & & & \\ 1 & 4 & 1 & & \\ & \ddots & \ddots & \ddots & \\ & & & 1 & 4 & 1 \\ & & & & 1 & 2 \end{bmatrix}$$

## 4. Fractional and integer order operational matrices of the integration for generalized hat functions

This section discussed the derivation [50] of operational matrices of fractional and integer order of the integration for generalized hat functions in the subsections 4.1 and 4.2 respectively.

### 4.1. Operational matrix of integer order integration of the generalized hat functions

Since  $\int_0^x \psi_i(x) dx \in L^2[0, T]$ , Eq. (6) is used to approximate it in the terms of the generalized hat basis functions as:

$$\int_0^x \psi_i(x) dx \cong \sum_{j=0}^n a_{ij} \psi_j(x) dx, \quad i = 0, 1, 2, \dots, n \quad (10)$$

The coefficients  $a_{ij}$  and according to Eq. (9) are given by

$$a_{ij} = \int_0^{jh} \psi_i(x) dx, \quad j = 0, 1, 2, \dots, n.$$

From Eqs. (8) and (10) we have

$$\int_0^x \Psi_{n+1} dx \cong P_{n+1} \Psi_{n+1}(x)$$

The coefficients  $a_{ij}$  will form a  $(n+1) \times (n+1)$  matrix  $P_{n+1}$  which is given by

$$P_{n+1} = \left(\frac{h}{2}\right) \begin{bmatrix} 0 & 1 & 1 & 1 & \cdots & 1 & 1 & 1 \\ 0 & 1 & 2 & 2 & \cdots & 2 & 2 & 2 \\ 0 & 0 & 1 & 2 & \cdots & 2 & 2 & 2 \\ 0 & 0 & 0 & 1 & \cdots & 2 & 2 & 2 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & \cdots & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 & \cdots & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & & 0 & 0 & 1 \end{bmatrix}$$

#### 4.2. Operational matrix of fractional order integration of the generalized hat functions

The fractional integration of order  $\alpha$  of the vector  $\psi_{n+1}(x)$  defined in (8) can be expressed as [50]:

$$(I^\alpha \Psi_{n+1})(x) = P_{n+1}^\alpha \Psi_{n+1}(x)$$

where the  $(n+1) \times (n+1)$  matrix  $P_{n+1}^\alpha$  is called the operational matrix of fractional integration [50,53]:

where

$$P_{n+1}^\alpha = \frac{h^\alpha}{\Gamma(\alpha+2)} \begin{bmatrix} 0 & \theta_1 & \theta_2 & \theta_3 & \cdots & \theta_n \\ 0 & 1 & \vartheta_1 & \vartheta_2 & \cdots & \vartheta_{n-1} \\ 0 & 0 & 1 & \vartheta_1 & \cdots & \vartheta_{n-2} \\ 0 & 0 & 0 & 1 & \cdots & \vartheta_{n-3} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

where  $\theta_k = k^\alpha(\alpha - k + 1) + (k - 1)^{\alpha+1}$ ,  $k = 1, 2, 3, \dots, n$ ; and  $\vartheta_k = (k + 1)^{\alpha+1} - 2k^{\alpha+1} + (k - 1)^{\alpha+1}$ ,  $k = 1, 2, 3, \dots, n - 1$ .

#### 5. The direct approach for solving fractional order variational problems using operational matrix of generalized hat basis function

In this section we shall consider the problem of extremization of a functional  $J$  of the form:

$$J[y(x)] = \int_0^1 F[x, y(x), {}_0^C D_x^\alpha y(x)] dx, \quad 0 < \alpha \leq 1 \quad (11)$$

Satisfying the condition  $y(0) = y_0$  and  $y(1)$  is considered to be undetermined.  ${}_0^C D_x^\alpha y$  refers to the fractional order derivative in the Caputo sense. The regular method for solving problem (11) is through the Euler-Lagrange equation, [54]:

$$\frac{\partial F}{\partial y} + {}_x^C D_1^\alpha \frac{\partial F}{\partial {}_0^C D_x^\alpha y} = 0$$

And

$$\left( \frac{\partial F}{\partial {}_0^C D_1^\alpha y} \right) \Big|_{x=1} = 0 \quad (12)$$

This section mainly uses the operational matrix of generalized hat functions to establish  $\psi_{n+1}$ . Now we approximate  ${}_0^C D_x^\alpha y$  in terms of hat basis functions as follows:

$${}_0^C D_x^\alpha y \approx C_{n+1}^T \Psi_{n+1} \quad (13)$$

where  $C_{n+1} = [c_0, c_1, c_2, \dots, c_n]^T$  and  $\Psi_{n+1}(x) = [\psi_0(x), \psi_1(x), \psi_2(x), \dots, \psi_n(x)]^T$ . Applying  $I_x^\alpha$  to the both sides of Eq. (13), hence  $y(x)$  can be expressed as:

$$y(x) \approx C_{n+1}^T P_{n+1}^\alpha \Psi_{n+1}(x) + y(0)$$

i.e.,

$$y(x) \approx C_{n+1}^T P_{n+1}^\alpha \Psi_{n+1}(x) + y_0 \quad (14)$$

The other terms in the functional of Eq. (11) are known functions of the independent variable  $x$  and can be expanded into hat basis functions through substitution, and finally we have:

$$J = J(c_0, c_1, c_2, \dots, c_n) \quad (15)$$

The original extremization of a fractional variational problem shown in Eq. (11) becomes the extremization of a functional of a finite set of variables in Eq. (15). Taking the partial derivative of  $J$  with respect to  $c_i$  and setting them equal to zero, we obtain

$$\frac{\partial J}{\partial c_i} = 0, \quad i = 0, 1, 2, \dots, n$$

Solving for  $c_i$ , and substituting into Eq. (14), one can reach to the desired solution of Eq. (11).

#### 6. Numerical examples

In order to have a comparison, a direct Haar wavelet method of Eq. (11) is briefly derived based on the previous work [44]. Assuming

$${}_0^C D_x^\alpha y(x) = \sum_{i=0}^{\infty} e_i h_i(x)$$

where  $h_i(x)$  indicates the Haar basis function [55]. Taking finite terms as an approximation, we have

$${}_0^C D_x^\alpha y(x) = \sum_{i=0}^{m-1} e_i h_i(x) = E_m^T H_m(x)$$

$$\text{Where } E_m^T = [e_0, e_1, e_2, \dots, e_{m-1}]^T \text{ and } H_m(x) = [h_0(x), h_1(x), h_2(x), \dots, h_{m-1}(x)]^T \quad (16)$$

Applying  $I_x^\alpha$  to the both sides of Eq. (16), thus  $y(x)$  can be written as

$$y(x) = E_m^T G_{m \times m}^\alpha H_m(x) + y_0 \quad (17)$$

where  $G_{m \times m}^\alpha$  is called the Haar wavelet operational matrix of fractional order integration [56]

Substituting (16) and (17) in the functional (11) and therefore we have

$$J = J(e_0, e_1, e_2, \dots, e_n)$$

Set  $\frac{\partial J}{\partial e_i} = 0$ ,  $i = 0, 1, \dots, m-1$ .

And solving for  $e_i$  and then substituting into (17) in order to get the desired solution.

**Example 1.** Consider the functional:

$$J[y(x)] = \int_0^1 \left[ \frac{1}{2} ({}_0^C D_x^\alpha y(x))^2 - y(x) \right] dx, \quad 0 < \alpha \leq 1 \quad (18)$$

and the boundary condition

$$y(0) = y_0 \text{ and } y(1) \text{ is unspecified} \quad (19)$$

First, we approximate  ${}_0^C D_x^\alpha y(x)$  in terms of hat basis functions as follows:

$${}_0^C D_x^\alpha y = C_{n+1}^T \Psi_{n+1}(x) \quad (20)$$

Here, we shall consider  $y_0 = 0$  and  $n = 8$  and therefore  $h = 0.125$ .

Now, upon taking  $I_x^\alpha$  to the both sides of Eq. (20) thus we get:

$$y(x) = C_{n+1}^T P_{n+1}^\alpha \Psi_{n+1}(x) \quad (21)$$

The other condition that we have is given by Eq. (12) and according to our example, we get:

$${}_0^C D_x^\alpha y(x)|_{x=1} = 0 \quad (22)$$

which implies that:

$$C_9^T \Psi_9(1) = 0$$

Therefore

$$C_8 = 0 \quad (23)$$

Substituting Eqs. (20), (21) and (23) into Eq. (18) we get:

$$J[y(x)] = \int_0^1 \left[ \frac{1}{2} (C_9^T \Psi_9(x) \Psi_9^T(x) C_9) - C_9^T P_9^\alpha \Psi_9(x) \right] dx$$

Hence:

$$J[y(x)] = \frac{1}{2} C_9^T \int_0^1 \Psi_9(x) \Psi_9^T(x) dx C_9 - C_9^T P_9^\alpha \int_0^1 \Psi_9(x) dx \quad (24)$$

Or

$$J[y(x)] = \frac{1}{2} C_9^T K_{9 \times 9} C_9 - C_9^T P_9^\alpha \begin{bmatrix} \frac{0.125}{2} \\ 0.125 \\ 0.125 \\ 0.125 \\ 0.125 \\ 0.125 \\ 0.125 \\ 0.125 \\ \frac{0.125}{2} \end{bmatrix}$$

Following Fig. 1 represent the approximate solution of Eq. (18) for different values of  $\alpha$  with a comparison with the exact solution when  $\alpha = 1$  which was given in [27] as:

$$y(x) = x(1 - \frac{x}{2})$$

Following Table 1 represent a comparison between the numerical solution of Eq. (18) using the proposed method and the Haar wavelet method with the exact solution when  $\alpha = 1$ .

**Example 2.** Find the extremal of the following functional:

$$J[y(x)] = \int_0^1 [({}_0^C D_x^\alpha y(x))^2 + x {}_0^C D_x^\alpha y(x)] dx, \quad 0 < \alpha \leq 1 \quad (25)$$

And the boundary condition

$y(0) = y_0$  and  $y(1)$  is unspecified

**Table 1**

Comparison between the numerical solution of Eq. (18) using the proposed method and Haar wavelet method with the exact solution when  $\alpha = 1$ .

$x$	The proposed method $n = 8$	Haar wavelet method $m = 8$	Exact solution
0	0	0.058	0
0.125	0.116	0.168	0.117
0.25	0.218	0.317	0.219
0.375	0.304	0.34	0.305
0.5	0.374	0.535	0.375
0.625	0.429	0.45	0.430
0.75	0.468	0.561	0.469
0.875	0.491	0.623	0.492

We also approximate  ${}_0^C D_x^\alpha y(x)$  in terms of hat basis functions with  $n = 8$ ,  $y_0 = 0$  and hence  $h = 0.125$  as follows:

$${}_0^C D_x^\alpha y = C_{n+1}^T \psi_{n+1}(x) \quad (26)$$

Hence:

$$y(x) = C_{n+1}^T P_{n+1}^\alpha \psi_{n+1}(x) \quad (27)$$

There is a variable  $x$  involved in Eq. (25) explicitly and it can be expanded in terms of hat basis function as:

$$x = d_{n+1}^T \psi_{n+1}(x) \quad (28)$$

where

$$d_{n+1} = [0 \quad 0.125 \quad 0.25 \quad 0.375 \quad 0.5 \quad 0.625 \quad 0.75 \quad 0.875 \quad 1]^T$$

By the transversality condition we have:

$$2 {}_0^C D_x^\alpha y(x) + x|_{x=1} = 0$$

Which implies that:

$$2 C_{n+1}^T \Psi_{n+1}(x) + x|_{x=1} = 0$$

$$C_{n+1}^T \Psi_{n+1}(1) = -\frac{1}{2}$$

Hence:


$$C_8 = -\frac{1}{2} \quad (29)$$

Substituting Eqs. (26), (28) and (29) into Eqs. (25) yields:

$$J[y(x)] \approx \int_0^1 [(C_9^T \Psi_9(x) \Psi_9^T(x) C_9) + C_9^T \Psi_9(x) \Psi_9^T(x) d_9] dx$$

Or

$$J[y(x)] = C_9^T K_{9 \times 9} C_9 + C_9^T K_{9 \times 9} d_9$$



**Figure 1.** Represent the numerical solution of Example (1) for different values of  $\alpha$  with the exact solution.




Figure 2. Represent the numerical solution of Example (2) for different values of  $\alpha$  with the exact solution.

Table 2

Comparison between the numerical solution of Eq. (25) using the proposed method and Haar wavelet method with the exact solution when  $\alpha = 1$ .

$x$	The proposed method $n = 8$	Haar wavelet method $m = 8$	Exact solution
0	0	0.001062	0
0.125	-0.003909	-0.008938	-0.003906
0.25	-0.016	-0.029	-0.016
0.375	-0.035	-0.049	-0.035
0.5	-0.063	-0.108	-0.063
0.625	-0.098	-0.120	-0.098
0.75	-0.141	-0.186	-0.141
0.875	-0.191	-0.222	-0.191

Following Fig. 2 represent the approximate solution of Eq. (25) for different values of  $\alpha$  with a comparison with the exact solution when  $\alpha = 1$  which was given in [57] as:

$$y(x) = -\frac{x^2}{4}$$

Following Table 2 represent a comparison between the numerical solution of Eq. (25) using the proposed method and the Haar wavelet method with the exact solution when  $\alpha = 1$ .

## 7. Conclusions

In this study a direct method based on operational matrix of generalized hat basis function was applied to solve fractional order variational problems. The fractional derivative is defined in the Caputo sense. The underlying problem was reduced under the proposed method to the problem of solving a set of algebraic equations which are solved by using an appropriate numerical technique.

The numerical solution for the class of problems considered can be obtained directly from the functional and there is no need to solve the fractional Euler-Lagrange equations.

By comparing the numerical solution using the proposed method with the exact solution and that of Haar wavelet method, we demonstrate the accuracy and efficiency of the proposed method for different values of  $\alpha$  even when  $n = 8$ .

## Acknowledgments

Author is very grateful to the reviewers for carefully reading the paper and for their comments and suggestions which have improved the paper.

## References

- [1] Sabatier J, Agrawal OP, Machado JAT. Advances in fractional calculus: theoretical developments and applications in physics and engineering. Springer; 2007.
- [2] Podlubny. Fractional differential equations. ACADEMIC press; 1999.
- [3] Momani S, Shawagfeh N. Decomposition method for solving fractional Riccati differential equations. Appl Math Comput 2006;182:1083–92.
- [4] Sweilam NH, Khader MM, Al-Bar RF. Numerical studies for a multi-order fractional differential equation. Phys Lett A 2007;371:26–33.
- [5] Tan Y, Abbasbandy S. Homotopy analysis method for quadratic Riccati differential equation. Commun Nonlinear Sci Num Simul 2008;13(3):539–46.
- [6] Khader MM. Introducing an efficient modification of the homotopy perturbation method by using Chebyshev polynomials. Arab J Math Sci 2012;18:61–71.
- [7] Bhrawy AH, Baleanu D, Assas L. Efficient generalized Laguerre-spectral methods for solving multi-term fractional differential equations on the half line. J Vib Control 2013;20:973–85.
- [8] Bhrawy AH, Doha EH, Ezz-Eldien SS, Gorder RAV. A new Jacobi spectral collocation method for solving 1+1 fractional Schrödinger equations and fractional coupled Schrödinger systems. Eur Phys J Plus 2014;129(12):1–21.
- [9] Heydari MH, Hooshmandasl MR, Mohammadi F, Cattani C. Wavelets method for solving systems of nonlinear singular fractional Volterra integro-differential equations. Commun Nonlinear Sci Num Simul 2014;19(1):37–48.
- [10] Ma J, Liu J, Zhou Z. Convergence analysis of moving finite element methods for space fractional differential equations. J Comput Appl Math 2014;255:661–70.
- [11] Bhrawy AH, Doha EH, Baleanu D, Ezz-Eldien SS. A spectral tau algorithm based on Jacobi operational matrix for numerical solution of time fractional diffusion-wave equations. J Comput Phys 2015;293:142–56.
- [12] Saadatmandi A, Dehghan M. A new operational matrix for solving fractional-order differential equations. Comput Math Appl 2010;59:1326–36.
- [13] Akrami M, Atabakzadeh M, Erjaee G. The operational matrix of fractional integration for shifted Legendre polynomials. Iran J Sci Technol 2013;37:439–44.
- [14] Bhrawy AH, Doha EH, Ezz-Eldien SS, Abdelkawy MA. A numerical technique based on the shifted Legendre polynomials for solving the time-fractional coupled KdV equation. Calcolo 2015. <http://dx.doi.org/10.1007/s10092-014-0132-x>.
- [15] Bhrawy AH, Zaky MA, Tenreiro Machado JA. Efficient Legendre spectral tau algorithm for solving two-sided space-time Caputo fractional advection-dispersion equation. J Vib Contr 2015. <http://dx.doi.org/10.1177/1077546314566835>.
- [16] Doha EH, Bhrawy AH, Ezz-Eldien SS. An efficient Legendre spectral tau matrix formulation for solving fractional sub-diffusion and reaction sub-diffusion equations. J Comput Nonlinear Dyn 2015;10(2):021019. 123.
- [17] Doha E, Bhrawy A, Ezz-Eldien S. A Chebyshev spectral method based on operational matrix for initial and boundary value problems of fractional order. Comput Math Appl 2011;62:2364–73.
- [18] Bhrawy A, Alofi A. The operational matrix of fractional integration for shifted Chebyshev polynomials. Appl Math Lett 2013;26:25–31.
- [19] Doha E, Bhrawy A, Ezz-Eldien S. A new Jacobi operational matrix: an application for solving fractional differential equations. Appl Math Model 2012;36:4931–43.
- [20] Doha EH, Bhrawy AH, Baleanu D, Ezz-Eldien SS. On shifted Jacobi spectral approximations for solving fractional differential equations. Appl Math Comput 2013;219:8042–56.
- [21] Kayedi-Bardeh A, Eslahchi M, Dehghan M. A method for obtaining the operational matrix of fractional Jacobi functions and applications. J Vib Control 2014;20:736–48.
- [22] Bhrawy A, Doha E, Baleanu D, Ezz-Eldien SS. A spectral tau algorithm based on Jacobi operational matrix for numerical solution of time fractional diffusion-wave equations. J Comput Phys 2015. <http://dx.doi.org/10.1016/j.icp.2014.03.039>.



- [23] Bhrawy A, Tharwat M, Alghamdi M. A new operational matrix of fractional integration for shifted Jacobi polynomials. *Bull Malays Math Sci Soc* 2014;37(4):983–95.
- [24] Bhrawy A, Baleanu D, Laila Assas, Efficient generalized Laguerre spectral methods for solving multi-term fractional differential equations on the half line. *J Vib Control* 2014;20:973–85.
- [25] Bhrawy AH. A Jacobi spectral collocation method for solving multi-dimensional nonlinear fractional sub-diffusion equations. *Num Algorithms* 2016. <http://dx.doi.org/10.1007/s11075-015-0087-2>.
- [26] Bhrawy AH, Taha TM. A review of operational matrices and spectral techniques for fractional calculus. *Nonlinear Dyn* 2016. <http://dx.doi.org/10.1007/s11071-015-2087-0>.
- [27] Agrawal OP. Formulation of Euler-Lagrange equations for fractional variational problems. *J Math Anal Appl* 2002;272(1):368–79.
- [28] El-Nabulsi RA, Torres DFM. Necessary optimality conditions for fractional action-like integrals of variational calculus with Riemann-Liouville derivatives of order  $(\alpha, \beta)$ . *Math Meth Appl Sci* 2007;30(15):1931–9.
- [29] Mozyrska D, Torres DFM. Modified optimal energy and initial memory of fractional continuous-time linear systems. *Signal Process* 2011;91(3).
- [30] Riewe F. Non concentrative Lagrangian and Hamiltonian mechanics. *Phys Rev E* 1996;53(2):1890–9.
- [31] Riewe F. Mechanics with fractional derivatives. *Phys Rev E* 1997;55(3):3582–92.
- [32] Agrawal OP. A new Lagrangian and a new Lagrange equation of motion for fractionally damped system. *J Appl Mech* 2001;68:339–41.
- [33] Klimek M. Fractional sequential mechanics-model with symmetric fractional derivatives. *Czechoslovak J Phys* 2001;51:1348–54.
- [34] Klimek M. Stationary conservation laws for fractional differential equations with variable coefficients. *J Phys A: Math Gen* 2002;35:6675–93.
- [35] Dreisigmeyer DW, Young PM. Extending Baures corollary to fractional derivatives. *J Phys A: Math Gen* 2004;37:L117–21.
- [36] Agrawal OP. A general formulation and solution scheme for fractional optimal control problems. *Nonlinear Dynam* 2004;38:323–37.
- [37] Tarasov VE, Zaslavsky GM. Fractional Ginzburg-Lunda equation for fractional media. *Phys A* 2005;354:249–61.
- [38] Agrawal OP. Fractional variational calculus in terms of Riesz fractional derivatives. *J Phys A* 2007;40:6287–303.
- [39] Agrawal OP. A general finite element formulation for fractional variational problems. *J Math Anal Appl* 2008;337:1–12.
- [40] Almeida R, Torres DFM. Calculus of variations with fractional derivatives and fractional integrals. *Appl Math Lett* 2009;22:1816–20.
- [41] Bastos N, Ferreira R, Torres DFM. Discrete-time fractional variational problems. *Signal Process* 2011;91:513–24.
- [42] Pooseh S, Almeida R, Torres DFM. A discrete time method to the first variation of fractional order variational functionals. *Cent Eur J Phys* 2013;11:1262–7.
- [43] Pooseh S, Almeida R, Torres DFM. Discrete direct methods in the fractional calculus of variations. *Comput Math Appl* 2013;66:668–76.
- [44] Osama HM, Fadhel SF, Zaid AM. Numerical solution of fractional variational problems using direct Haar Wavelet method. *Int J Innovative Res Sci Eng Technol* 2014;3(5).
- [45] Bhrawy AH, Doha EH, Tenreiro Machado JA, Ezz-Eldien SS. An efficient numerical scheme for solving multi-dimensional fractional optimal control problems with a quadratic performance index. *Asian J Control* 2015;17(6):2389–402.
- [46] Bhrawy AH, Doha EH, Baleanu D, Ezz-Eldien SS, Abdelkawy MA. An Accurate numerical technique for solving fractional optimal control problems. In: *Proceedings of the Romanian Academy, Series A*, vol. 16, Number 1; 2015. p. 47–54.
- [47] Bhrawy AH, Ezz-Eldien SS. A new Legendre operational technique for delay fractional optimal control problems. *Calcolo*. [http://dx.doi.org/10.1007/s10092-015-0160-1\(2016\)](http://dx.doi.org/10.1007/s10092-015-0160-1(2016)).
- [48] Ezz-Eldien SS, Hafez RM, Bhrawy AH, Baleanu D, El-Kalaawy AA. New numerical approach for fractional variational problems using shifted Legendre orthonormal polynomials. *J Optim Theory Appl* 2016. <http://dx.doi.org/10.1007/s10957-016-0886-1>.
- [49] Babolian E, Mordad M. A numerical method for solving systems of linear and nonlinear integral equations of the second kind by hat basis function. *Comput Math Appl* 2011;62:187–98.
- [50] Tripathi MP, Baranwal VK, Pandey RK, Singh OP. A new numerical algorithm to solve fractional differential equations based on operational matrix of generalized hat functions. *Commun Nonlinear Sci Num Simulat* 2013;18:1327–40.
- [51] Heydari MH, Hooshmandasl MR, Ghaini FMM, Cattani C. A computational method for solving stochastic it<sup>o</sup>-volterra integral equations based on stochastic operational matrix for generalized hat basis functions. *J Comput Phys* 2014;270:402–15.
- [52] Heydari MH, Hooshmandasl MR, Cattani C, Ghaini FMM. An efficient computational method for solving nonlinear stochastic it<sup>o</sup> integral equations, Application for stochastic problems in physics. *J Comput Phys* 2015;283:148–68.
- [53] Heydari MH, Hooshmandasl MR, Shakiba A, Cattani C. An efficient computational method based on the hat function for solving fractional optimal control problems. *Tbilisi Math J* 2016;9(1):143–57.
- [54] Agrawal OP. Fractional variational calculus and the transversality conditions. *J Phys A: Math Gen* 2006;39:10375–89.
- [55] Chen C, Hsiao C. Haar wavelet method for solving lumped and distributed-parameter systems. *IEEE Proc-Control Theory Appl* 1997;144(1):87–94.
- [56] Zhao W, Li Y. Haar wavelet operational matrix of fractional order integration and its applications in solving the fractional order differential equation. *Appl Math Comput* 2010;216(8):2276–85.
- [57] Hsiao CH. Haar wavelet direct method for solving variational problems. *Math Comput Simul* 2004;64:569–85.